

BBA 4257

Glaser ✓

EFFECTS OF HIGH-FREQUENCY ELECTRIC FIELDS ON THE LIVING CELL

I. BEHAVIOUR OF HUMAN ERYTHROCYTES IN HIGH-FREQUENCY ELECTRIC FIELDS AND ITS RELATION TO THEIR AGE

A. A. FÜREDI AND I. OHAD

Department of Biological Chemistry, The Hebrew University of Jerusalem, Jerusalem (Israel)

(Received July 22nd, 1963)

SUMMARY

Behaviour of human erythrocytes in high-frequency electric fields is described. When the field is applied erythrocytes show a reversible elongation which is accompanied by a rotatory motion. Old erythrocytes do not behave in this manner but form chains oriented in the direction of the field. The results can be explained by the interaction of two effects: (1) the polarization of the erythrocyte and (2) the distortion of the applied field by the fixed structural charges of the erythrocyte.

The method of separating young from old erythrocytes according to their specific gravity has been improved.

INTRODUCTION

Previous investigations have shown that microscopic particles become arranged parallel to the direction of electromagnetic fields in the radio-frequency range^{1,2}. It was demonstrated that this phenomenon is the result of the induced dipole moment of the particle³.

In the present work the behaviour of human erythrocytes in the high-frequency field has been investigated using the apparatus described before³.

MATERIALS AND METHODS

Erythrocytes were obtained from heparinised fresh blood. After centrifugation at $2500 \times g$ for 5 min the cells were resuspended in aqueous sucrose solution (11%, w/v, for isotonic and 5%, w/v, for hypotonic solutions) and stored for the duration of the experiment (1-3 h) at room temperature.

Specimens for examination were prepared by applying a drop of the cell suspension on a microscope cover glass (22 × 22 mm) which was then covered by a second round cover glass (19 mm diameter) and placed on the electrodes under the phase microscope. Specimens were usually allowed to stand 1-2 min before the field was applied for 1-5 sec from a balance oscillator of 120 megacycles/sec. Longer exposure (> 10 sec) causes damage to cells by heating.

A simple and quick procedure developed for separating between young and old erythrocytes, was based on the method of PRANKERD⁴. Erythrocytes were sedimented as described above and the plasma was discarded. One volume (0.2 ml) of packed cells was mixed gently with two volumes (0.4 ml) of 30% (w/v) crystalline bovine albumin (Armour Pharmaceutical Co. Ltd.) in a 5 × 50 mm plastic tube (Spinco Co.). The mixture was centrifuged in the swinging-bucket rotor of the Clay-Adams serological centrifuge at 3000 rev./min for 30 min. Glass serological centrifuge tubes were used as adaptors for the small plastic tubes.




Fig. 1. Tube with separated erythrocytes. Top layer, young erythrocytes; middle layer, albumin solution; bottom, old erythrocytes.

Centrifugation causes the cells to concentrate into two distinct layers, with clear and sharp boundaries (Fig. 1), a top layer of about 12 mm and a bottom one of 2-3 mm, separated by about 20 mm of albumin. The top layer was found to consist of young cells and the bottom layer of old cells when examined by the methods and criteria of DANON AND MARIKOVSKY¹⁰. During the process of fractionation some hemoglobin is released to the albumin solution. To separate between layers the tube is cut into two in the middle of the albumin layer. The cells from each layer are washed in 2 ml isotonic sucrose and saline (1:1 by volume) and centrifuged as described above. The washed packed cells (0.08 ml) were mixed with 1 ml of sucrose (isotonic or hypotonic) and 0.04 ml saline for further examination in the electric field.

RESULTS

When the field is applied to erythrocytes suspended in isotonic sucrose solution they become oriented so that their radii are perpendicular to the direction of the field

Fig. 2. Schematic




Fig. 3. Erythrocyt




Fig. 4. Erythrocyt




Fig. 2. Schematic drawing of the amoeba-like rotation of erythrocytes suspended in isotonic sucrose solution when an electric field is applied.




Fig. 3. Erythrocytes suspended in isotonic sucrose solution oriented 45° to the field direction (\rightarrow).




Fig. 4. Erythrocytes suspended in hypotonic sucrose solution elongated parallel to the field direction (\rightarrow).

and old
limented
of packed
ie bovine
inco Co.).
y-Adams
uge tubes

tyer, albumin

rs, with clear
ottom one of
nd to consist
methods and
i some hemo-
ie tube is cut
r, are washed
as described
rose (isotonic
field.

solution they
n of the field

a, 79 (1964) 1-8

and rotate slowly. The rotation of the cells results from distortions of shape reminiscent of amoeboid movement (Fig. 2). The cell undergoes this change of shape in such a way that Point 1 (Fig. 2a) remains static while the upper part of the cell (A) increases in volume on account of the lower part (B). When the field is turned off while the shape of the erythrocyte is still distorted (Fig. 2b, c or d) it returns to its natural form (Fig. 2a or 2e). If the field is cut off when the cell has reached position e (Fig. 2) no further change in shape takes place, indicating that the cell has reached a stable form.

In addition to rotation, during a prolonged exposure to the electric field (5 sec) cells form long chains oriented about 45° to the direction of the field (Fig. 3).

Cells suspended in hypotonic sucrose solution lose their biconcave form and swell to spherical shape. When electric field is applied they elongate so that their long axis is parallel to the direction of the field (Fig. 4) and rotate without changing its direction (Fig. 5). Elongation of erythrocyte is 2-4 times greater than the normal




Fig. 5. Schematic representation of the effect of field on young erythrocytes in hypotonic medium. t_0 represents undistorted erythrocyte at zero time. t_1 , t_2 show distortion and rotation of erythrocyte. Point A describes an ellipse around central axis. Cross hatching represents the orientation of the cell contents.

cell diameter. Addition of salt ($\geq 0.02\%$ NaCl) stops cell rotation but does not prevent elongation. If the elongation is maintained for 5-15 sec, cells break up into small spherical vesicles which sometimes remain interconnected by long thin filaments. The process of vesicle formation seems to be preceded by a contact between opposite regions of cell membranes at points of maximum elongation. The opacity of the vesicles appears to be similar to that of whole cells indicating that there is no extensive leakage of the cell contents. Morphologically, they resemble the myelinic forms described by BESSIS¹¹.

Finally, prolonged application of the electric field results in excessive heat formation, cells break up and release their contents. The remaining ghosts are no longer affected by the field. These ghosts are granulated and differ in appearance from those obtained by gradual hemolysis⁶ and which *do react* to the electric field similarly to intact cells.

Not all the cells in blood samples obtained from healthy donors became elongated under hypotonic conditions. This difference in behaviour may be explained by differences in membrane properties of the individual cells. Separating the cells into top (young) and bottom (old) fractions using the procedure described, it was found that more than 90% of the top fraction but less than 10% of the bottom fraction exhibited rotation and elongation (Fig. 6a, b). Old erythrocytes often form chains oriented in the direction of the field.





Fig. 6. Influence of field

shape in such
cell (A) im-
med off while
to its natural
ne (Fig. 2) no
stable form.
c field (5 sec)
Fig. 3).
hal form and
so that their
out changing
in the normal

atomic medium.
tion of erythro-
the orientation

s not prevent
up into small
in filaments.
ween opposite
of the vesicles
ensive leakage
was described

excessive heat
ghosts are no
in appearance
e electric field

time elongated
explained by
the cells into
l, it was found
ation fraction
n form diam-




Fig. 6a.




Fig. 6b.

Fig. 6. Influence of field on erythrocytes separated into old and young fractions. a, top fraction (young); b, bottom fraction (old).

DISCUSSION

The various types of behaviour of erythrocytes in the high-frequency electric field can be explained in terms of their dielectric constant, structural charges, shape, elasticity and composition of the external medium.

The described behaviour of erythrocytes can be better understood if we first consider model systems: uncharged particles with a dielectric constant different from that of the surrounding medium which are polarized in a high-frequency electromagnetic field and form long chains parallel to the direction of the field³. If these particles were elongated they would align themselves with their long axis parallel to the lines of force (Fig. 7a). On the other hand, elongated particles with fixed structural charges align themselves at right angles to the applied field^{7,8}. As a result of the distortion of the field, elongated particles would tend to assume a resting position perpendicular to lines of force where the field distortion and hence the potential energy is minimal (Fig. 7b). The degree of field distortion resulting from structural charges on the particle and the intensity of its polarisation mutually determine the final orientation. This prediction has been verified experimentally with bacteria which are charged elongated particles (B. HERSKOVITS, unpublished experiments). Raising the salt concentration or lowering the pH of the medium surrounding the bacteria causes them to reorient in the direction of the field. This is readily explained since the fixed charges are neutralised by added ions while the induced dipole is not affected³.

On the basis of the above considerations, an attempt has been made to predict the behaviour of idealised particles in a high-frequency electric field. The influence of the field on rigid and elastic charged and uncharged spheres and ellipsoids is described in Table I.




Fig. 7. Schematic drawing of distortion of electric field by particles and the resulting stresses. Stress (P) resulting from field distortion according to MAXWELL⁷ causes elongation: a. Elastic uncharged particle of low dielectric constant. Stress (P) causes elongation and orientation parallel to the field (→). b. Rigid charged particle. Stress (P') causes orientation perpendicular to field direction (←→).

The elongation (Fig. 5) corresponds to the data in Table I. This is due to the fact that the discs with a low dielectric constant are two conflicting forces: they tend to align themselves in the direction of the field. This last alignment must be in the manner shown in Fig. 5.

Young erythrocytes in a characteristic manner in terms of the isotonic and hypotonic solutions, resulting in a characteristic shape.

The addition of salt thus inhibits rotation (Table I).

Old erythrocytes rotate, but can form long chains (Table I) and young cells do not rotate. The dielectric constant while the field is applied. Different chemical properties work two other properties are described in Table I.

THEORETICAL BEHAVIOUR FROM

| Spheres | Rigid |
|------------|---------|
| | Elastic |
| Ellipsoids | Rigid |
| | Elastic |

* Chain-type orientation

The elongation and rotation of young erythrocytes under hypotonic conditions (Fig. 5) corresponds closely to the behaviour predicted for elastic charged spheres (Table I). This could be explained if it is assumed that these cells are charged elastic discs with a dielectric constant lower than that of the medium. Apparently there are two conflicting influences on the erythrocyte: polarisation would tend to elongate them in the direction of the field, but as they bear fixed structural charges they would tend to align themselves perpendicular to the field, like charged elongated particles. This last alignment is not possible since the elongation which results from polarisation must be in the field direction. The overall result is that the particles rotate in the manner shown in Fig. 5.

Young erythrocytes in isotonic solution do not elongate and rotate more slowly in a characteristic manner (Fig. 2). This behaviour is too complicated to explain in terms of the idealized systems (Table I) but may be partly due to a higher charge density on the cell surface. The difference in behaviour between erythrocytes in isotonic and hypotonic solutions could be due to a decrease in charge density in the latter, resulting from swelling.

The addition of salt to young erythrocytes reduces the effect of their charge, thus inhibits rotation and they exhibit the behaviour of uncharged elastic spheres (Table I).

Old erythrocytes in both hypotonic and isotonic solution do not elongate or rotate, but can form chains oriented parallel to the field direction like rigid uncharged spheres (Table I). A possible explanation of the difference in behaviour between old and young cells is that decreasing lipid content during aging⁴ increases their dielectric constant while their charge density decreases⁹ and hence they become less sensitive to the field. Differentiation between young and old erythrocytes has been based on chemical properties, radioactive tagging⁴ and on morphological studies⁵. In the present work two other possible methods for the quantitation of old and young cells in blood samples are described. Firstly, the modification of the method of PRANKERD combined

TABLE I

THEORETICAL BEHAVIOUR OF SPHERES AND ELLIPSOIDS OF DIELECTRIC CONSTANT DIFFERENT FROM THAT OF THE MEDIUM IN A HIGH-FREQUENCY ELECTRIC FIELD

The addition of salt transforms charged particle to uncharged.

| | | | Change in shape | Rotary motion | Orientation in the field direction | Orientation perpendicular to the field | Chain formation | Roulaux |
|------------|---------|-----------|--------------------|------------------|--|--|--------------------|---------|
| Spheres | Rigid | Charged | - | - | - | - | - | + |
| | | Uncharged | - | - | + | - | + | - |
| | Elastic | Charged | + | + | - | + | - | - |
| | | Uncharged | + | - | + | - | + | - |
| Ellipsoids | Rigid | Charged | - | - | - | + | - | + |
| | | Uncharged | - | - | + | - | + | - |
| | Elastic | Charged | + | - | - | + | - | - |
| | | Uncharged | + | - | + | - | + | - |

* Chain-type orientation.


ency electric field
charges, shape,

od if we first con-
different from that
y electromagnetic

If these particles
parallel to the lines
structural charges
of the distortion
ion perpendicular
energy is minimal
charges on the
final orientation.

which are charged
ng the salt concentra-
causes them to
the fixed charges
ed³.

made to predict
ld. The influence
and ellipsoids is



ie resulting stresses.
ngation. a. Elastic
ion and orientation
tation perpendicular

with the estimation of haemoglobin in the two fractions, could be conveniently used clinically. Secondly, the difference in behaviour of young and old cells in the electric fields combined with photomicrography could be applied to differential counting of the two types of cells (elongated *vs.* round).

The possibility exists that under more carefully controlled conditions, examination of erythrocytes in high-frequency electric fields could be applied to the study and detection of pathological conditions.

ACKNOWLEDGEMENTS

Our thanks are due to Professor M. SCHRAMM, Professor D. DANON and Mr. R. ZILBERBERG for criticism and suggestions during this work. We are also thankful to Dr. D. M. SERR and Mrs. B. SHINKOLEVSKY for providing us blood samples and to Miss A. AVINESER for technical assistance. This work was supported by a grant from the Hirschman Foundation, in Memory of Dr. A. SHAMGAR and in part by Research Grant D-2 from the Ford Foundation through the Israel Foundation Trustees.

REFERENCES

- ¹ A. A. TEIXEIRA-PINTO, L. L. NEJELSKI, J. L. CUTLER AND J. H. HELLER, *Exptl. Cell Res.*, 20 (1960) 548.
- ² E. MUTH, *Kolloid-Z.*, 41 (1927) 97.
- ³ A. A. FÜREDI AND R. C. VALENTINE, *Biochim. Biophys. Acta*, 56 (1962) 33.
- ⁴ T. A. PRANKERD, *J. Physiol.*, 143 (1958) 325.
- ⁵ D. DANON AND K. PERK, *J. Cellular Comp. Physiol.*, 59 (1962) 117.
- ⁶ D. DANON, A. NEVO AND Y. MARKOVSKY, *Bull. Res. Council Israel*, 6E (1956) 36.
- ⁷ C. A. COULSON, *Electricity*, Oliver and Boyd, Edinburgh-London, 1961, p. 64.
- ⁸ C. J. F. BÖTTCHER, *Theory of Electric Polarisation*, Elsevier, Amsterdam, 1952, p. 62.
- ⁹ D. DANON AND Y. MARIKOVSKY, *Compt. Rend.*, 253 (1961) 1271.
- ¹⁰ D. DANON AND Y. MARIKOVSKY, *Compt. Rend. Soc. Biol.*, Tome CLV No. 1 (1961) 12.
- ¹¹ M. BESSIS, *Traité de Cytologie Sanguine*, Masson, Paris, 1954.

Biochim. Biophys. Acta, 79 (1964) 1-8

BBA 4247

FLUC

JEROME

Depa

The fluorescence is lower than in terms of the surface and of the induction. The fluorescence of the induction compared with the energy transfer a result of photoreduction intensifies to a lower level rises to a new deficiency, the slowly to the temporary increase in photosynthesis.

Variations with a valuable tool in literature and concentrations experimental data that two different re-investigations two different fields described the sequence an earlier report

* Present address: Turkey.
** Present affiliation: Chemistry, University